

How to take

LAGEVRIO[®] (molnupiravir)


This medicine has been given to you to treat your COVID-19 disease.

LAGEVRIO[®] has provisional approval for the treatment of adults with COVID-19 who do not require initiation of oxygen due to COVID-19 and who are at increased risk for hospitalisation or death. The decision to approve this indication is based on the efficacy and safety data from a Phase 3 trial. Continued approval of this indication depends on additional data.

How to take

Take LAGEVRIO® exactly as your doctor tells you.


Take 4 capsules immediately when you receive your medicine.




Take 4 capsules of LAGEVRIO® every 12 hours (for example, in the morning at 9 am and in the evening at 9 pm).



Take LAGEVRIO® for 5 days.



Take LAGEVRIO® with a full glass of water, with or without food.



Do not open, break, or crush the capsules. If you cannot swallow capsules whole, tell your healthcare provider.

Pregnancy, contraception and breastfeeding

Pregnancy

LAGEVRIO® is not recommended in pregnancy. If there is a chance you might be pregnant, talk to your doctor about having a pregnancy test prior to commencing LAGEVRIO®.

Contraception

If you are female:

Use effective birth control while you are taking LAGEVRIO® and for 4 days after stopping if there is a possibility of you getting pregnant.

If you are male:

If you are a male who is sexually active with a partner who has the potential to become pregnant, use a reliable method of contraception during treatment and for 3 months after the last dose of LAGEVRIO®.

Breastfeeding

Breastfeeding is not recommended during treatment and for 4 days after the last dose of LAGEVRIO®.

Talk to your doctor if you have any other medical conditions, take any other medicines, or are pregnant or plan to become pregnant or are breastfeeding or plan to breastfeed.



What to do if you miss a dose?



If it has been **less than 10 hours** since the missed dose, take it as soon as you remember.



If it has been **more than 10 hours** since the missed dose, skip the missed dose and take your dose at the next scheduled time.

Do not take a double dose to make up for the dose you missed. Call your doctor or pharmacist if you are not sure what to do.

LAGEVRIO[®] Dosing Calendar

LAGEVRIO[®] should be taken regularly at the same times each day.


It is important that you do not miss or skip doses of this medicine.



Take 4 capsules every 12 hours, with a full glass of water, with or without food, for 5 days.

Fill in the table on the right - and cross out each day as you take them, to help you keep track of your doses.

If your first dose of LAGEVRIO[®] is on the evening of Day 1, then you will need to take your last dose of LAGEVRIO[®] on the morning of Day 6.



MORNING DOSE TAKEN AT: _____ AM

DAY 1	DAY 2	DAY 3	DAY 4	DAY 5
DATE / /	DATE / /	DATE / /	DATE / /	DATE / /

EVENING DOSE TAKEN AT: _____ PM

DAY 1	DAY 2	DAY 3	DAY 4	DAY 5
DATE / /	DATE / /	DATE / /	DATE / /	DATE / /

For more information

For more information, including any side effects you may experience, please read the full Consumer Medicine Information (CMI). The CMI is available by visiting this link: **msdinfo.com.au/lagevriocmi**.

Please contact your doctor or pharmacist if you have any questions.



▼ This medicine is subject to additional monitoring. This will allow quick identification of new safety information. You can help by reporting any side effects you may get. You can report side effects to your doctor, or directly at www.tga.gov.au/reporting-problems.